

¹Laboratory of Molecular Carcinogenesis, MD C4-06, National Institute of Environmental Health Sciences, National Institutes of Health, Research Triangle Park, North Carolina, USA.

²Section of Molecular Carcinogenesis, Institute of Cancer Research, Haddow Laboratories, Surrey, UK and Chester Beatty Laboratories, Fulham Road, London, UK

³Department of Molecular Medicine, Endocrine Tumor Unit, Karolinska Hospital, Stockholm, Sweden

⁴Department of Human Genetics and Medicine, McGill University, Montreal General Hospital, Montreal, Quebec, Canada

⁵Department of Surgery, Duke University Medical Center, Durham, North Carolina 27710-2611, USA

⁶CRC Center for Molecular Biology, Institute for Cancer Research, Haddow Laboratories, Surrey, UK and Chester Beatty Laboratories, Fulham Road, London, UK

⁷Department of Obstetrics and Gynecology, Division of Gynecologic Oncology, Duke University Medical Center, Durham, North Carolina, USA

⁸Department of Genetics, Duke University Medical Center, Durham, North Carolina, USA

Correspondence should be addressed to P.A.F. e-mail: futre001@mc.duke.edu

BRCA2 mutations in primary breast and ovarian cancers

Johnathan M. Lancaster¹, Richard Wooster², Jonathon Mangion², Catherine M. Phelan^{3,4,5}, Charles Cochran¹, Curtis Gumbs⁵, Sheila Seal², Rita Barfoot², Nadine Collins², Graham Bignell², Sandeep Patel², Rifat Hamoudi², Catharina Larsson³, Roger W. Wiseman¹, Andrew Berchuck⁷, J. Dirk Iglehart⁵, Jeffrey R. Marks⁵, Alan Ashworth⁶, Michael R. Stratton² & P. Andrew Futreal^{5,7,8}

The second hereditary breast cancer gene, *BRCA2*, was recently isolated¹. Germline mutations of this gene predispose carriers to breast cancer, and, to a lesser extent, ovarian cancer. Loss of heterozygosity (LOH) at the *BRCA2* locus has been observed in 30–40% of sporadic breast and ovarian tumours, implying that *BRCA2* may act as a tumour suppressor gene in a proportion of sporadic cases^{2–5}. To define the role of *BRCA2* in sporadic breast and ovarian cancer, we screened the entire gene for mutations using a combination of techniques in 70 primary breast carcinomas and in 55 primary epithelial ovarian carcinomas. Our analysis revealed alterations in 2/70 breast tumours and none of the ovarian carcinomas. One alteration found in the breast cancers was a 2-basepair (bp) deletion (4710delAG) which was subsequently shown to be a germline mutation, the other was a somatic missense mutation (Asp3095Glu) of unknown significance. Our results suggest that *BRCA2* is a very infrequent target for somatic inactivation in breast and ovarian carcinomas, similar to the results obtained for *BRCA1*.

We analysed genomic DNA from 70 breast and 55 epithelial ovarian cancers for *BRCA2* mutations. Thirty-four of the breast and 18 of the ovarian tumours (56% and 53% respectively of informative cases) showed LOH in the *BRCA2* region, using the markers *D13S260*, *D13S171*, *D13S260*, and *D13S153*. The latter marker is intragenic to the *RB1* gene. All tumours analysed which were informative for both the *BRCA2* and the *RB1* regions of 13q show concomitant LOH

Table 1 *BRCA2* mutations in primary breast cancer

Patient no.	Age	Mutation	Effect	Germline/somatic
19	59	4710delAG	frameshift	germline
6	74	C to A	Asp3095Glu	somatic

(data not shown). The complete coding region was screened in each sample using both single strand conformation analysis (SSCA) and denaturing gel deletion analysis. In addition, exons 10, 11 and 27 were also screened by the protein truncation test (PTT). Potential sequence alterations represented by aberrant bands were characterized by direct sequencing in all cases.

Sequence alterations were identified in 2/70 breast tumours (Table 1), but none of the 55 ovarian samples. An SSCA shift in exon 11 was detected in breast carcinoma sample #19 (Fig. 1). This fragment, along with the corresponding fragment from the patient's normal lymphocyte DNA and a healthy control, were sequenced directly. A 2-bp AG deletion was detected, corresponding to a 4710delAG mutation. This deletion produces a frameshift leading to a predicted premature termination codon 2 amino acids downstream. The deletion was present in the patient's germline (Fig. 1b). No family history of cancer had been reported by this individual, who was diagnosed at age 59 with a unilateral infiltrating ductal carcinoma. In breast carcinoma sample #6 a somatic alteration was detected. This alteration was found to be a C to A transversion in exon 25 resulting in an aspartate to glutamate amino acid change at codon 3095 (Fig. 2). The change was not present in the patient's lymphocyte DNA, nor in over 300 control chromosomes. Six different highly polymorphic microsatellite markers on chromosomes 16 and 17 confirmed that the tumour and lymphocyte DNA came from the same patient (data not shown). This tumour sample had LOH in the *BRCA2* region and it was the aberrant glutamate residue that was retained in the tumour. This patient had a unilateral infiltrating ductal adenocarcinoma diagnosed at age 74. The significance of this amino acid change is unclear. Additionally, several sequence variants were detected via SSCA⁶.

The lack of *BRCA2* mutations in sporadic breast and ovarian cancers is very reminiscent of the results obtained for *BRCA1*^{7–10}. Similar to the observation for *BRCA1*, the region containing *BRCA2* undergoes LOH in a fraction of breast and ovarian carcinomas⁸. The




Fig. 1 a, SSCA autoradiogram showing shift in Patient 19 in lane 3. b, ABI electropherograms showing germline delAG in patient 19. The antisense strand is shown. MC19T, DNA from patient's tumour. MC19N, DNA from patient's peripheral blood. Control is unrelated normal blood DNA.




Fig. 2 a, Autoradiogram SSCA showing shift of tumour 6 in lane 3. b, Electropherogram showing sequence of missense mutation in tumour 6. Antisense strand is shown for both patients' tumour and normal blood DNA. The G to T transversion in tumour DNA shown in the right panel results in a GAC to GAA/ Asp to Glu amino acid substitution on the sense strand. This missense change was not present in the matching peripheral blood DNA from the same patient in the left panel.

Table 2a BRCA2 PCR primers

Exon	Primers	Sequence (5'→3')	Exon	Primers	Sequence (5'→3')
2	JAD46	F-CAAGCATGGAGGAATATCG R-GTTTTATGGTCTTAAGCAACAC	JAD8		AACCAGAAAGAATAAATACT CCTCAACGCAAAATCTTCAT
3	JAD47	CACAATTTGTCTGTCACTGG ATTGCATTAICTACTAAGTC	JAD7		TTCCAAGTAATATCCAATGTA ATTTTTGATTTATCTCGTGTIT
4	JAD48	ATCCAGAGTATATACATTTCTC GATCTTCTACCAGGGCTCTTAG	JAD6		AAGTGAAGACATATTTACAGACAG TATGAAGCTTCCCTATACT
5,6	JAD 52	AAAATAACCTAAGGGATTTGC CAAATCTCAATTAATAAGTC	JAD5		CACCTTGTGATGTTAGTTTG TTGGGATTAATAATGTTCTGGAGTA
7	JAD49	CAATTCAGTAAACGTTAAGTG AACAGAAAGTATTAGAGATGAC	JAD4		AAGTAAACGAACTCAGACC CTGGGTTTCTCTTATCAACAGCA
8	JAD50	CATGTAATCAAAATAGTAGATGTC CTCAAAGGCTTAGATAAATTACAG	JAD3		GTCTTCACTATTCACTACAG AGTGAGACTTTGGTCTTAAT
9	JAD51	GCATTGAGAGTTTTATACTAGTG ACCTGTAGTCAACTAAACAG	JAD2		TTCAACAAGCAAAACAACAGT GTGAGTTCATCATCTTCCATAAA
10	JAD54F JAD23R	GCCTCTGTTTTATACATTTAAACAGG GATCAGTATCATTTGGTCTCCAC	JAD1		CTTACTCCAAAGATTCAGAAAATC AGCATACCAAGTCTACTGAATAAC
	JAD20F JAD23R	CAAAAGCCACATGGAAAGTC GATCAGTATCATTTGGTCTCCAC	12	JAD30	ATAAAAGTATGATTTATTTGCC TCCACCTGAGGTCAGTAAT
	JAD24	AAGCAACGCTGATGAATGTG TGGTCAACATGAAGAAATATG	13	S1237 S1236	TAAAGCCATAATTTGTCTCA CTTCTTAAAGCTTAAAGTGCATT
	JAD25	CAGGTCTAATGAGGCCACAG GAGAAGTCCAGATATTGCC	14	A45 A44	AAGGAACGTCAGAGATACAG GGTTGGTCTGCCTGTAGTAAT
	JAD26	CATCTTGAATCTCATACAGAC GACATAAGGAGTCTCCTCTTC		S1239 S1238	CTTCAAGCAATTTAGCAGTTTCAGG GOTGCTTGAATGGAGTTGTT
	JAD27F JAD20R	AAGCCCTCTGAAAGTGGACTG GCCAATGTAAGTGGTCTTC	15	JAD31	ATTTAATTACAAGTCTTCAG ACTCTGTCTATAAAAAGCCATC
11	JAD53	GATGGTACTTAAATTTGTCAAC CAAGATCCCTGAGAGTACTG	16	JAD32	TATTTTGTGAGCTGTATACG AGGGAATCATAAAAAGTTAAC
	JAD22	GCTCTTTGGGCAAACTCTG ATAAAAGACTTTTCTGGGATTG	17	JAD33	ATTCAGTATCATCTTATGGT TATGATTACGTAATGTAATGC
	JAD21	TGGAATACAGTACTACTGAC TTTTCAAGTGGCAACAGCTC	18	JAD34	GAATCTAGAGTCAACCTCC ATCTAAGTGGCCCTTAACAGC
	JAD19	CCCATTGAAAAGATCAAGATG GTTCTTAGTATTCCTAAAGC	19	JAD35	TGGCCATTATGAACTTACAG AATGAGCATCTTATGTAAGC
	JAD18	TGCTTCCAAAGTAGCTAATG CTGTGATTTGAAATGGACC	19	JAD36	GAATGAAAACCTTATGATATCTG AAGAGACCAGAAATTTAAAGAGC
	JAD17	ACATGAACAATGGGACAGAC TGGTTTGAATTAATCCTGTC	20	JAD37	CACGTGCTGGCCCTGATAC AGTCTTAAAGACTTTGTTCTC
	JAD16	GTCATATAACCCCTCAGATG CTGTACCTTCAAATGGCTTGC	21	JAD53	TAAATCTCCCTCTTTGGGTG TTCCCTTCTGTAGTGGCCAG
	JAD15	CGATTGGTCAGGTAGACAGC CTCTGAGAAAGTTTCTCTCAC	22	JAD38	TCTAGTTACAATAGATGGAAC AATCATTTTGTAGTAAAGTC
	JAD14	TGTTTCTACTGAAGCTCTGC GTTATCTCATTTTCAATTTCTC	23	JAD39	GCATCTTCTCATCTTTCTCC TGAAATAAAATTCATCTGAAAAC
	JUL5 F3 JUL5 R4	TTGAAATGACTACTGGCCAC CCTTCATAAACTGGCCAGATAAT	24	JAD40	TTGTTAGTTTGAAGTCTCC TAATCATAAAGAGATTTAAAGAGC
	JAD13	TGCTTAAATATCTGGCCAC AAATGACTCTTTGGCGACAC	25	JAD41	TTCCATTCTAGGACTTGGCC GTGGTGAATGCTGAAAAGTAAAC
	JAD12	AGATTTTGGAGCTTCTGATAC TCCAGTCAACCACTGGGACAC	26	JAD42	TTTATAAGCAGCTTTTCCAC ATACCTTCTATAATTTCTGTAG
	JUL5 F1 JUL5 R1	TGGACATCTAAGTTATGAGG ATTTCACCTAGTACCCTTCTTTT	27	JAD43	ACATAATTATGATAGGCTACG AAATGTACAATGGGACTAAC
	JAD11	TGATGAAAAAGAGCAGGATG ACAAGGTTTATCATTTATG	JAD44		AGCCTTGGATTTCTTGAGTGT TCTAGTGGATTCACCTGACAG
	JAD10	CTGCCCAAGGTGTAAGAAAT AATGACTGAATAGGGGACTGAT	JAD45		TCTTTTGTCTGGTCAACAGG AAGCGTCAATAATTTATGTG
	JAD9	TCCTGCAACTTGTACAC GATTTTGTCTTTTCAGC			

relatively close proximity of the *RB1* gene could be driving the majority of the LOH seen in these cases. Our study set contained no tumours that demonstrated loss at one region versus the other when informative for both. However, for breast and particularly ovarian cancers, there is evidence to suggest that the *RB1* gene can be excluded from the regions of loss in some cases and is not always targeted for inactivation in cases with large scale losses on 13q (refs 3, 5, 11). Our screening set contained both tumours selected for LOH in the *BRCA2* region as well as unselected cancer cases to increase the likelihood of finding mutations. Additionally, we employed three complementary techniques in screening for mutations, and thus feel that few coding region mutations have been missed. The two alterations found were both from tumours showing LOH. One case was a germline frameshift deletion which was retained in the tumour, the other a somatic missense mutation which was also retained in the tumour. The significance of the latter somatic mutation is unclear. Our data showing infrequent mutations in sporadic tumours suggests that if *BRCA2* plays a significant role in tumorigenesis in the non-hereditary forms of these cancers, it is through a mechanism other than structural mutation. Further, combined with the data on somatic mutations of *BRCA1* in sporadic breast and ovarian cancers, the evidence suggests that hereditary breast cancer (and ovarian cancer in the context of breast/ovarian cancer syndromes) may be fundamentally different diseases at the molecular genetic level. This is not to suggest that these genes play no role in the development of non-hereditary breast and ovarian cancers. Recent evidence would suggest that subcellular localization and/or expression levels may be critical in *BRCA1* involvement in cancer^{12,13}. Given the results presented here, parallel studies on the expression, localization and, ultimately, normal function of *BRCA2* are paramount.

Methods

Samples. Tumour tissue and matched blood lymphocytes were obtained from patients treated at Duke University Medical Center, USA, and the Royal Marsden Hospital, England and from the Gynecologic Oncology Group/Cooperative Human Tissue network ovarian tissue bank (USA). Tissues were obtained under general consent for discarded tissue and tumour/white cell banking. Genomic DNA was obtained from tumour tissue and blood using standard procedures. The breast cancers were all infiltrating ductal carcinomas with the exception of two pure intraductal carcinomas. The mean age of onset for the breast study set was 53. The ovarian carcinomas were of mixed histology, the majority being papillary serous (90%), and the mean age of onset for the tumours studied was 58. There were no cases of either bilateral breast or dual primary breast/ovarian cancer in the study set.

Single strand conformation analysis. The entire coding region, including intron/exon borders, was examined by SSCA (using primers given in Table 2a). Genomic DNA (20 ng) was amplified using primers under the following standard PCR conditions: 50 mM KCl, 10 mM Tris-HCl, pH 9.0, 1.5 mM MgCl₂ (Promega), 0.2 mM dTTP, dCTP and dGTP, 0.05 mM dATP, and (α-³²P)dATP (Amersham) at 1 μCi/reaction and 2 U *Taq* DNA Polymerase (Promega) in a final volume of 10 μl. Thermocycling conditions consisted of 30 cycles of 30 s at 94 °C, 30 s at 55 °C, and 30 s at 72 °C, followed by one cycle of 3 min extension at 72 °C in a 9600 Thermocycler (Perkin-

Table 2b BRCA2 PTT primers

Exon	Primers	PTT primer sequence ^a (5'→3')
10	JAD20F × JAD20R	JL-PTTH GAAACAGTTGTAGATACCTCTGAAGA
11	JAD54F × JAD17R	JL-PTTD GATTCTGAAGAACCACTTTGTCC
11	JAD17F × JAD14R	JL-PTTF GAAATCAAGCTCTCTGAACATAAC
11	JAD14F × JAD12R	JL-PTTE GAACTTCTGCAGAGGTACATCCA
11	JAD12F × JAD8R	JL-PTTA GACATTCTAAGTTATGAGGA
11	JAD9F × JAD5R	JL-PTTB GGTCACCAGAAAGAATAAATACT
11	JAD5F × JAD2R	JL-PTTG GGGAAAGCTTCATAAGTCAGTC
27	JAD43F × JAD45R	JL-PTTJ TCTTCTCCTAATTGTGAGATA

^aEach PTT primer is preceded by the T7/Kozak sequence 5'-GGATCCTAATACGACTCACTATAGGGAGACCACCATG-3'.

Elmer). PCR product (4 µl) was diluted in 56 µl of loading buffer (95% formamide, 10 mM NaOH, 0.05% bromophenol blue, 0.05% xylene cyanol), denatured at 94 °C for 5 min, and rapidly cooled on ice. A sample of 4 µl was electrophoresed through a 0.5× MDE gel (AT Biochem) containing 0.6× TBE buffer, at room temperature for 14–18 h at 8 W, dried, and autoradiographed for 1–18 h. Migration shift analysis was also used for detection of small deletions/insertions¹. Samples were amplified as described above and were run through 5% denaturing sequencing gels at 70 W, dried and autoradiographed for 1–4 h.

Protein truncation test. Exons 10, 11 and 27, which represent 64% of the coding region, were amplified in 8 segments by the polymerase chain reaction (PCR) using primers given in Table 2b. A 1 µl aliquot of each of these primary templates was subjected to 10 additional cycles of PCR, in a reaction mixture containing a modified primer carrying a T7 promoter and eukaryotic translation initiation sequence in place of the forward primer. A 20 µl reaction mix containing 6 µl of the secondary PCR product, 1.6 µl of ³⁵S-methionine (1,000

Ci/mmol, Amersham), 0.8 µl RNasin (recombinant 40 U/µl), 0.4 µl TnT T7 RNA Polymerase, 0.2 µl amino acid Methionine(-) Mix, and 5.0 µl TnT Rabbit reticulocyte lysate (Promega), was incubated at 30 °C for 1 h. This product (6 µl) was electrophoresed on a 10–20% SDS-polyacrylamide Ready-Gel (Biorad), fixed, dried and autoradiographed for 3–18 h.

DNA sequence analysis. Sequencing templates were produced for samples showing aberrant mobility on SSCA or PTT. Aberrant SSCA bands were cut from the MDE gel and eluted in 100 µl of dH₂O for 90 min at 37 °C. An additional 30 cycles of PCR was then carried out to amplify the DNA eluted from the gel slice. A parallel PCR was carried out to amplify genomic DNA from the same patient using the same primers. PCR products were purified using the Wizard PCR Prep DNA Purification System (Promega) and sequenced using a PRISM DyeDeoxy Terminator Cycle Sequencing kit and a 373 automated fluorescent sequencer (Applied BioSystems), according to manufacturer's instructions.

Acknowledgements

We thank P. Biggs, N. Rahman and B. Gusterson for assistance. This work was funded in part by the NCI/Duke University Specialized Program of Research Excellence (SPORE) in breast cancer P50-CA68438, NIH/NCI grant R21-CA68348, NIH grant 1R21-CA66228, the National Institutes of Environmental Health Sciences, the Cancer Research Campaign, BREAKTHROUGH Breast Cancer Charity and Jean Rook Appeal, the US Army, the Institute for Cancer Research. C.M.P. and C.L. were funded by the Swedish Cancer Foundation.

Received 28 February; accepted 29 March 1996.

1. Wooster, R. *et al.* Identification of the breast cancer susceptibility gene *BRCA2*. *Nature* **378**, 789–792 (1995).
2. Lundberg, C. *et al.* Loss of heterozygosity in human ductal breast tumours indicates a recessive mutation on chromosome 13. *Proc. Natl. Acad. Sci. USA* **84**, 2372–2376 (1987).
3. Kim, T.M. *et al.* Loss of heterozygosity on chromosome 13 is common only in the biologically more aggressive subtypes of ovarian epithelial tumors and is associated with normal retinoblastoma expression. *Cancer Res.* **54**, 605–609 (1994).
4. Collins, N. *et al.* Consistent loss of the wild type allele in breast cancers from a family linked to the *BRCA2* gene chromosome 13q12–13. *Oncogene* **10**, 1673–1675 (1995).
5. Cleiton-Jansen, A.M. *et al.* Loss of heterozygosity in sporadic breast tumors at the *BRCA2* locus on chromosome 13q12–13. *Br. J. Cancer* **72**, 1241–1244 (1995).
6. Phelan, C.M. *et al.* Mutation analysis of the *BRCA2* gene in 49 site-specific breast cancer families. *Nature Genet.* **13**, 120–122 (1996).
7. Futreal, P.A. *et al.* *BRCA1* mutations in primary breast and ovarian carcinomas. *Science* **266**, 120–122 (1994).
8. Merajver, S.A. *et al.* Somatic mutations in the *BRCA1* gene in sporadic ovarian tumours. *Nature Genet.* **9**, 439–443 (1995).
9. Hosking, L. *et al.* A somatic *BRCA1* mutation in an ovarian tumour. *Nature Genet.* **9**, 343–344 (1995).
10. Takahashi, H. *et al.* Mutation analysis of the *BRCA1* gene in ovarian cancers. *Cancer Res.* **55**, 2998–3002 (1995).
11. Kerangueven, F. *et al.* Patterns of loss of heterozygosity at loci from chromosome arm 13q suggest a possible involvement of *BRCA2* in sporadic breast tumors. *Genes Chrom. Cancer* **13**, 291–294 (1995).
12. Chen, Y. *et al.* Aberrant subcellular localization of *BRCA1* in breast cancer. *Science* **270**, 789–791, (1995).
13. Thompson, M.E., Jensen, R.A., Obermiller, P.S., Page, D.L. & Holt, J.T. Decreased expression of *BRCA1* accelerates growth and is often present during sporadic breast cancer progression. *Nature Genet.* **9**, 444–450 (1995).